

ANALYSIS OF GEOMAGNETIC STORM VARIATIONS AND COUNT-RATE OF COSMIC RAY MUONS RECORDED AT THE BRAZILIAN SOUTHERN SPACE OBSERVATORY

Everton Frigo¹, Jairo Francisco Savian², Marlos Rockenbach da Silva³, Alisson Dal Lago⁴,
Nalin Babulal Trivedi⁵ and Nelson Jorge Schuch⁶

Recebido em 23 janeiro, 2006 / Aceito em 29 setembro, 2006
Received on January 23, 2006 / Accepted on September 29, 2006

ABSTRACT. An analysis of geomagnetic storm variations and the count rate of cosmic ray muons recorded at the Brazilian Southern Space Observatory – OES/CRS/INPE-MCT, in São Martinho da Serra, RS during the month of November 2004, is presented in this paper. The geomagnetic measurements are done by a three component low noise fluxgate magnetometer and the count rates of cosmic ray muons are recorded by a Muon Scintillator Telescope – MST, both instruments installed at the Observatory. The fluxgate magnetometer measures variations in the three orthogonal components of Earth magnetic field, H (North-South), D (East-West) and Z (Vertical), with data sampling rate of 0.5 Hz. The Muon Scintillator Telescope records hourly count rates. The arrival of a solar disturbance can be identified by observing the decrease in the muon count rate. The goal of this work is to describe the physical morphology and phenomenology observed during the geomagnetic storm of November 2004, using the H component of the geomagnetic field and vertical channel V of the multi-directional muon detector in South of Brazil.

Keywords: geomagnetic storms, geomagnetic variations, cosmic rays, interactions Earth-Sun, space weather.

RESUMO. Neste trabalho apresentamos uma análise das variações geomagnéticas e contagem de raios cósmicos (muons) detectados no Observatório Espacial do Sul – OES/CRS/INPE – MCT (29.43°S, 53.80°W), em São Martinho da Serra, RS, durante o mês de Novembro de 2004. As medidas das variações geomagnéticas são realizadas através de um magnetômetro *fluxgate* de três componentes e baixo ruído, e as contagens de muons são obtidas através de um detector de muons multi-direcional, ambos instrumentos instalados no Observatório. O magnetômetro *fluxgate* detecta variações nas três componentes ortogonais do campo magnético Terrestre, H (Norte-Sul), D (Leste-Oeste) e Z (Vertical), com uma taxa de aquisição de 0.5 Hz. O Telescópio Cintilador de Muons realiza contagens horárias. A chegada de um distúrbio solar é identificada observando-se o decréscimo na contagem de muons. O objetivo deste trabalho é fazer uma descrição da morfologia e fenomenologia física observada durante a tempestade geomagnética de Novembro de 2004, usando a componente H do campo geomagnético e o canal vertical V do Telescópio Cintilador de Muons no Sul do Brasil.

Palavras-chave: tempestades, variações geomagnéticas, raios cósmicos, interações Terra-Sol, clima espacial.

¹University of São Paulo, USP, Institute of Astronomy, Geophysics and Atmospheric Sciences, IAG/USP, Department of Geophysics, Rua do Matão, 1.226, Cidade Universitária – 05508-090 São Paulo, SP, Brazil. Phone: +55 (11) 3091-2789; Fax: +55 (11) 3091-5034 – E-mail: efrigo@iag.usp.br

²Space Science Laboratory of Santa Maria, LACESM/CT, UFSM Southern Regional Space Research Center, CRS/INPE, MCT Campus of the Federal University of Santa Maria, UFSM, Center of Technology, LACESM, Cidade Universitária, P.O. Box 5021 – 97105-900 Santa Maria, RS, Brazil. Phone: +55 (55) 3220-8021; Fax: +55 (55) 3220-8007 – E-mail: savian@lacesm.ufsm.br

³National Institute for Space Research, INPE/MCT, Division of Space Geophysics, DGE, Av. dos Astronautas, 1.758, P.O. Box 515 – 12245-970 São José dos Campos, SP, Brazil. Phone: +55 (12) 3945-6754; Fax: +55 (12) 3945-6810 – E-mail: marlos@dge.inpe.br

⁴National Institute for Space Research, INPE/MCT, Division of Space Geophysics, DGE, Av. dos Astronautas, 1.758, P.O. Box 515 – 12245-970 São José dos Campos, SP, Brazil. Phone: +55 (12) 3945-6768; Fax: +55 (12) 3945-6979 – E-mail: dallago@dge.inpe.br

⁵Partnership FATEC/UFSM, FURNAS, INPE, National Institute for Space Research, INPE/MCT, Division of Space Geophysics, DGE, Av. dos Astronautas, 1.758, P.O. Box 515 – 12245-970 São José dos Campos, SP, Brazil. Phone: +55 (12) 3945-6807; Fax: +55 (12) 3945-6810 – E-mail: trivedi@dge.inpe.br

⁶Southern Regional Space Research Center, CRS/INPE, MCT, Campus of the Federal University of Santa Maria, UFSM, Center of Technology, LACESM, Cidade Universitária, P.O. Box 5021 – 97105-900 Santa Maria, RS, Brazil. Phone: +55 (55) 3220-8021; Fax: +55 (55) 3220-8007 – E-mail: njschuch@lacesm.ufsm.br

INTRODUCTION

Areas of instability in the Sun can release high-speed plasma structures, with great amounts of matter and energy, the so-called coronal mass ejections – CMEs (Hundhausen et al., 1984; Hundhausen, 1997) throughout the whole 11-year solar cycle. Although more frequent during maximum phase, these CMEs are also present during low activity periods of the solar cycle. Eventually these solar CMEs reach the Earth (Schwenn et al., 2005), interacting with the earth's magnetosphere. A portion of the incoming solar plasma may enter into the magnetosphere via a reconnection process (Dungey, 1961), causing great disturbances in the Earth's magnetic field denominated geomagnetic storms (Gonzalez et al., 1994), observed by ground magnetic observatories. These geomagnetic storms can be classified according to different Dst index levels: weak: $-50 \text{ nT} \leq \text{Dst} \leq -30 \text{ nT}$; moderate $100 \text{ nT} \leq \text{Dst} \leq -50 \text{ nT}$ and intense $\text{Dst} < -100 \text{ nT}$ (Gonzalez et al., 1994). This index is based on hourly measures of the H component of the geomagnetic field at selected low latitude stations. The principal characteristic of a geomagnetic storm is the decrease of the H component of the geomagnetic field (Kamide et al., 1998). This decrease is attributed to the increase of the population of particles trapped in magnetosphere. The solar structures capable of causing geomagnetic storms move through the solar wind shielding the passage of galactic cosmic rays, causing decreases in their count rates on the ground, also known as "Forbush decreases" (Cane, 1993; Cane et al., 1994, 1996). In this work we analyzed the variations observed in the H component of the geomagnetic field and the decrease in the high energy cosmic ray muon count rates during the geomagnetic storm of November 2004, both observed at the Southern Space Observatory, in São Martinho da Serra, RS.

DATA AND METHODOLOGY OF ANALYSIS

The Brazilian Southern Space Observatory is located in the proximities of the center of South Atlantic Magnetic Anomaly – SAMA, that is, the area where the smallest intensity of the Earth's magnetic field is observed in the surface of the Earth (Trivedi et al., 2005). A low noise fluxgate magnetometer was used to measure the three orthogonal components of the geomagnetic field and the data were recorded at a sampling rate of 2 seconds (0.5 Hz). We used the data of the variations observed in the H component during the period from 5 to 15 November, 2004, in order to identify the geomagnetic storm. The Muon Scintillator Telescope located at the Observatory, in South of Brazil, is able to observe high energy cosmic rays (muons) in several directions (Da Silva et al., 2004), and it is part of the International Muon Detector Network, with other ones at Nagoya (Japan) and Hobart (Australia). It re-

ports hourly count rates of the muon precipitation on the surface of the Earth. As described previously, a relationship exists among the solar disturbances, the variations in the geomagnetic field and the decrease in the muons count rate.

RESULTS

The Dst index is presented in Figure 1, obtained from Kyoto University (Japan), for November 2004. On November 8th, 2004, a decrease is observed in the Dst index, reaching a negative peak of -373 nT , indicating the occurrence of an intense geomagnetic storm. On November 10th, 2004 another decrease is observed, reaching -289 nT , indicating the occurrence of a second geomagnetic storm. Figure 2, shows the variations observed in the H component of the geomagnetic field for the period of 5-15 November 2004, detected by the fluxgate magnetometer. Figure 2 also shows the counts for the same period, detected by the muon-scintillator telescope at the Southern Space Observatory. The vertical line 1 of Figure 2 indicates an accentuated decrease of approximately 3,5% in the muon count rate, indicating that a Coronal Mass Ejection (CME) was driven towards the Earth, reaching the earth's magnetosphere. Line 2 displays a sudden increase in the intensity of the H component and a subsequent decrease, indicating that a solar structure impact the Earth magnetosphere. Line 3 displays a second decrease in the muon count rate, of approximately 4%, indicating that another solar structure reached the Earth. Line 4 shows a sudden increase in the intensity of the H component, and subsequent decrease, indicating that the second solar structure impact the Earth's magnetosphere.

CONCLUSIONS

Based on these results, it is observed that the Muon Scintillator Telescope Network may detect the passage of solar structures (CMEs) by the Earth's magnetosphere. It is of high importance for Space Weather to detect the arrival of shocks and storms, from detailed studies of these structures. In the near future, cosmic ray data will be used to detect the arrival of CMEs before they reach the earth, using data from the whole high energy cosmic ray network. It is essential to detect them with a large antecedence, in order to avoid eventual damages, such as: intensification of electric currents in space and at the Earth surface, occurrence of polar auroras, acceleration of charged particles, and several damages in satellites, damages in the Global Positioning System (GPS), in telecommunications and even to the astronauts that are in spaceships.




Figure 1 – Dst index of the month of November 2004.




Figure 2 – Variations in H Component of Earth's Magnetic Field and Count Rate of the Muon Scintillator Telescope for the period of 5-15 November 2004.

ACKNOWLEDGEMENTS

Thanks to World Data Center for Geomagnetism, Kyoto University, for the Dst index, and to the International Solar-Terrestrial

Physics Project. We thank the group of research workers at the Space Weather Laboratory, to Shinshu University for technical support and to the CNPq for the financial support: PIBIC/Process 104441/2004-6 and 107615/2003-7, and partial support from

project 472396/04-8 and also to Fundação de Amparo à Pesquisa do Estado de São Paulo – FAPESP for the financial support by the project 05/54800-1.

REFERENCES

- CANE HV. 1993. Cosmic ray decreases and magnetic clouds. *J. Geophys. Res.*, 98: 3509–3512.
- CANE HV, RICHARDSON IG & von ROSENINGE TT. 1996. Cosmic ray decreases: 1964-1994. *J. Geophys. Res.*, 101(A10): 21561-21572.
- CANE HV, RICHARDSON IG, von ROSENINGE TT & WIBBERENZ G. 1994. Cosmic Ray Decreases and Shock Structure: A Multispacecraft Study. *J. Geophys. Res.*, 99: 21429–21441.
- DA SILVA MR, CONTREIRA DB, MONTEIRO S, TRIVEDI NB, MUNAKATA K, KUWABARA T & SCHUCH NJ. 2004. Cosmic ray muon observation at southern space observatory – SSO (29°S, 53°W). *Astrophysics and Space Science*, 290: 389–397.
- DUNGEY JW. 1961. Interplanetary magnetic field and the auroral zones. *Phys. Rev. Lett.*, 6(2): 47–48, January.
- GONZALEZ WD, JOSELYN JA, KAMIDE Y, KROEHL HW, ROSTOKER G, TSURUTANI BT & VASYLIUNAS VM. 1994. What is a magnetic storm? *J. Geophys. Res.*, 99(A4): 5771–5792, April.
- HUNDHAUSEN AJ. 1997. An Introduction. In: CROOKER N, JOSELYN JA & FEYNMAN J. (Ed.). *Coronal mass ejections*, Washington, DC: AGU, 99: 1–7.
- HUNDHAUSEN AJ, SAWYER CB, HOUSE LL, ILLING RME & WAGNER WJ. 1984. Coronal mass ejections observed during the solar maximum mission – latitude distribution and rate of occurrence. *J. Geophys. Res.*, 89(A5): 2639–2646.
- KAMIDE Y, YOKOYAMA N, GONZALEZ WD, TSURUTANI BT, DAGLIS IA, BREKKE A & MASUDA S. 1998. Two-Step development of geomagnetic storms. *J. Geophys. Res.*, 103(A4): 6917–6921, April.
- SCHWENN R, DAL LAGO A, HUTTUNEN E, GONZALEZ WD. 2005. The association of coronal mass ejection with their effects near the Earth. *Ann. Geophys.*, 23: 1033–1059.
- TRIVEDI NB, PATHAN BM, SCHUCH NJ, BARRETO LM & DUTRA SLG. 2005. Geomagnetic Phenomena in the South Atlantic Anomaly Region in Brazil. *Advances in Space Research*, England, 36: 2021–2024.

NOTES ABOUT THE AUTHORS

Everton Frigo is Undergraduate in Physics (2005) at the Federal University of Santa Maria – UFSM, Santa Maria, Brazil. He developed during three years (2003-2005) activities of Undergraduate Research, studying the relationships between geomagnetic variations observed at the Earth's surface and the Earth-Sun interaction, at the Laboratory of Solar Physics, Interplanetary Medium and Magnetospheres at the Southern Regional Space Research Center of the National Institute for Space Research – CRS/INPE – MCT, in Santa Maria. Currently, he is M.Sc. Student in Geophysics at the Institute of Astronomy, Geophysics and Atmospheric Sciences – IAG in the University of São Paulo – USP, São Paulo, Brazil.

Jairo Francisco Savian is a Physics Undergraduate student, attending the ninth semester, at the Federal University of Santa Maria – UFSM, trainee at the National Institute for Space Research – INPE. Currently he is developing his fourth year of science initiation research at the Southern Regional Space Research Center, Space Weather Laboratory – SWL/CRS/INPE – MCT, in Santa Maria. He is working with the data analysis of the Multidirectional Muon Detector Telescope on the Project “Analysis of the solar and interplanetary origins of intense geomagnetic storms”, in cooperation with the Space Science Laboratory of Santa Maria – LACESM/CT – UFSM.

Martos Rockenbach da Silva is Undergraduate in Physics at the Federal University of Santa Maria – UFSM in 2003. He got his M.Sc. Degree on Space Geophysics in 2005 at the National Institute for Space Research, São José dos Campos, Brazil. Currently he is Ph.D. student at the National Institute for Space Research, in São José dos Campos.

Alisson Dal Lago is Undergraduate in Physics (1996) at the Federal University of Santa Maria – UFSM, Brazil, Master (1999) and Ph.D. (2003) in Space Geophysics at the National Institute for Space Research – INPE, Brazil, currently research scientist (since 09/2004) of the Space Geophysics Division – DGE, Atmospheric and Space Sciences – CEA of INPE, where he works with topics related to Space Weather, study and forecasting, using ground and space observations.

Nalin Babulal Trivedi is Undergraduate in Physics (1960), M.Sc. (1962), and Ph.D. (1969), at the University of Gujarat in Ahmedabad, India. He is researcher (since 1970) in the area of geomagnetism at the Space Geophysics Division – DGE, Atmospheric and Space Sciences – CEA, in the National Institute for Space Research – INPE, Brazil.

Nelson Jorge Schuch is Undergraduate in Physics at the Federal University of Santa Maria – UFSM, in 1972. M.Sc. (Physics) – Extra Galactic Astrophysics, University of Mackenzie, Brazil, in 1975. Ph.D. (Astrophysics) – University of Cambridge, England, in 1979. Post Doctoral Experience at Cambridge University, in 1979/1980. From 1980 to 1995, he worked as Vice-Director of the National Observatory – ON/MCT, Brazil. He is the Head of the Southern Regional Space Research Center, CRS/INPE/MCT. His research fields are the Astrophysics, with emphasis in Radioastronomy/Observational Cosmology, and also the Space and Atmospheric Sciences, with emphasis in the Earth-Sun Interactions, Space Weather, Aeronomy, Space Geophysics.